

RIGENOMA®

CREAM

Repairs and Regenerates the Skin

with



OZOILE®
ERBAGIL PATENT
Stable Ozonides

INDICATIONS

RIGENOMA® CREAM with Ozoile® is indicated for prevention and treatment of acute and chronic skin lesions with different etiology also in case of risk of infection such as bedsores, vascular, metabolic, pressure ulcers, surgical wounds, traumatic injuries and from chemo and radiotherapy. The product is also indicated in the prevention and treatment of dermatological pathologies such as Acne, Psoriasis, Dermatitis and Dermatoses with different etiology such as Atopic dermatitis, Eczematous dermatitis, all characterized by inflammation, irritation, erythema, xerosis, itching, desquamation, possible infection, alterations of the flora and skin barrier.

MAIN COMPONENTS: Ozoile® (Stable Ozonides with Vitamin E acetate), Ruscus Aculeatus Root Extract, Calendula Officinalis Flower Extract, Ribes Nigrum Leaf Extract, Equisetum Arvense Extract, Sodium Hyaluronate, Thiocctic Acid.

Ozoile®, Stable ozonides with Vitamin E acetate, is a pool of innovative molecules obtained from a Patented process that has anti-inflammatory and anti-itching action, counteracts hypoxia, creates an unfavorable environment to pathogenic microbial proliferation and promotes tissue repair processes.

HOW TO USE

Spread the product directly on the affected area. In the treatment of acute and chronic skin lesions with different etiology, after cleansing with Rigenoma® detergent and subsequent application of Rigenoma® spray on the bottom of lesion, apply a homogeneous layer of Rigenoma® cream on the peri-wound skin and apply a veil to the surrounding area for more than 5 cm. The product can also be applied directly to the entire

injured area especially during the granulation and re-epithelialization phase.

Rigenoma® cream can be used in combination with Rigenoma gauze® to cover the lesion to protect, keep the environment moist and to promote tissue repair and regeneration by actively working with the base of the wound. Apply Rigenoma® cream once or twice a day or in case of bandage and occlusive dressing every three or four days to renew the dressing as indicated by physician. In the treatment of dermatological problems, apply the product 2 or 3 times a day. The treatment period differs according to the rapidity of therapeutic response.

WARNINGS AND PRECAUTIONS

Topical use. Do not ingest. Keep out of the reach of children. Do not use it if the packaging appears to be damaged. Do not use the device beyond the expiration date. Expiration date refers to the intact and correctly stored product. Store in a cool, dry place away from heat sources and direct sunlight. Store preferably between 5 °C and 35 °C. Wash your hands well before using the medical device. Carefully close the tube after use. After opening it is recommended to use the product within 6 months.

CONTRAINdications

Rigenoma® cream is well tolerated. It is contraindicated in case of ascertained hypersensitivity towards the components. There are no cases of sensitization or side effects if this phenomenon occurs please discontinue use and consult your physician.

There are no known overdoses cases.

There are no known interactions with drugs and medicinal substances.

PACKAGE

100 ml - 40 ml tube.

GLUTEN FREE

NO COLOURS

NO ALLERGENS

 ERBAGIL s.r.l.
Via L. Settembrini, 13
82037 Telesio Terme (BN) Italy
www.erbagil.com

REV. 03 02/2024

  0425



RIGENOMA®

KREMA

Obnavlja in regenerira kožo

S
sestavino



OZOILE®
ERBAGIL PATENT
Stabilni ozonidi

INDIKACIJE

RIGENOMA® KREMA s sestavino Ozoile® je indicirana za preprečevanje in zdravljenje akutnih in kroničnih lezij z različno etiologijo, tudi v primeru nevarnosti okužbe, kot so preležanine, vaskularne in presnovne razjede, razjede zaradi pritiska, kirurške rane, poškodbe in lezije zaradi kemoterapije in radioterapije.

Izdelek je indiciran tudi za preprečevanje in zdravljenje dermatoloških bolezni, kot so akne, luskavica, dermatitis in dermatozra z različno etiologijo, kot sta atopijski dermatitis in ekcematozni dermatitis, za katere so značilni vnetje, draženje, rdečina, kseroza, srbenje, luščenje, možnost okužbe ter spremembe flore in kožne pregrade.

GLAVNE SESTAVINE: Ozoile® (stabilni ozonidi z vitaminom E acetatom), izvleček iz korenin bodeče lobodičke (Ruscus aculeatus), izvleček iz cvetov vrtnega ognjiča (Calendula officinalis), izvleček iz listov črnega ribeza (Ribes nigrum), izvleček njivske preslice (Equisetum arvense), natrijev hialuronat, alfa-lipoična (tioktična) kislina.

Ozoile®, stabilni ozonidi z vitaminom E acetatom, je skupke inovativnih molekul, pridobljenih s patentiranim postopkom, ki deluje protivnetno in proti srbenju, preprečuje hipoksijo, ustvarja neugodno okolje za razmnoževanje patoloških mikrobov in pospešuje procese popravila tkiva.

KAKO SE UPORABLJA

Izdelek namažite neposredno na prizadeto površino. Pri zdravljenju akutnih in kroničnih kožnih lezij z različno etiologijo najprej očistite z Rigenoma® čistilnim sredstvom in dno lezije popršite z Rigenoma® pršilom. Nanesite enakomerno plast Rigenoma® krema na kožo okoli rane in prekrijte vsaj 5 cm okolice s kontaktno mrežico.

Izdelek se lahko nanese tudi neposredno na celotno poškodovanov površino, zlasti med fazama granulacije in ponovne epitelizacije. Rigenoma® krema se lahko uporablja skupaj z Rigenoma® gazo za pokrivanje lezije za zaščito, ohranjanje vlažnega okolja in za pospeševanje popravila in obnove tkiva z aktivnim delovanjem na dno rane. Nanesite Rigenoma® kremo enkrat ali dvakrat na dan oziroma v primeru povijanja in okluzijske obloge vsake tri ali štiri dni po navodilih zdravnika, da obnovite oblogo.

Pri zdravljenju kožnih težav nanesite izdelek 2 ali 3-krat na dan. Trajanje zdravljenja se razlikuje glede na hitrost terapevtskega odziva.

OPOZORILA IN PREVIDNOSTNI UKREPI

Topikalna uporaba.

Ne zaužijte.

Shranjujte izven dosega otrok.

Ne uporabljajte, če je embalaža na videz poškodovana.

Pripomočka ne uporabljajte po datumu izteka roka uporabnosti.

Rok uporabnosti se nanaša na nedotaknjen in pravilno shranjen izdelek.

Shranjujte na hladnem, suhem mestu, stran od virov topote in neposredno sončne svetlobe.

Shranjujte po možnosti pri temperaturi med 5 °C in 35 °C. Pred uporabo medicinskega pripomočka si dobro umijte roke.

Po uporabi tubo skrbno zaprite.

Po odprtju je priporočljivo uporabiti izdelek v 6 mesecih.

KONTRAINDIKACIJE

Krema Rigenoma® se dobro prenaša.

Kontraindicirana je v primeru potrjene preobčutljivosti na sestavine.

Ni znanih primerov preobčutljivosti ali stranskih učinkov. Če pride to tega pojava, prenehajte z uporabo in se posvetujte z zdravnikom.

Ni znanih primerov prevelikega odmerjanja.

Ni znanih primerov medsebojnega delovanja z zdravili in zdravilnimi snovmi.

PAKIRANJE

Tuba s 100 ml ali 40 ml.

BREZ GLUTENA

BREZ BARVIL

BREZ ALERGENOV

Distributer:
MagPharm,
prodaja farmacevtskih izdelkov, d.o.o.
Litostroška 44e,
1000 LJUBLJANA

ERBAGIL s.r.l
Via L. Settembrini, 13
82037 Telesse (BN) Italy
www.erbagil.com

REV. 03/2024

MD C E 0425

